



Optimizing the control of foot-and-mouth disease in Denmark by simulation Meta-analysis on the efficacy of foot-and-mouth disease emergency vaccination

Hisham Beshara Halasa, Tariq; Boklund, Anette ; Enøe, Claes

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Hisham Beshara Halasa, T. (Author), Boklund, A. (Author), & Enøe, C. (Author). (2012). Optimizing the control of foot-and-mouth disease in Denmark by simulation: Meta-analysis on the efficacy of foot-and-mouth disease emergency vaccination. Sound/Visual production (digital)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Optimizing the control of foot-and-mouth disease in Denmark by simulation

Meta-analysis on the efficacy of foot-and-mouth disease emergency vaccination


Tariq Halasa

Anette Boklund

Claes Enøe

DTU Vet

National Veterinary Institute



Background

- EU member state must maintain the FMD free status. In case of an outbreak, animals within a defined area could be vaccinated.
- To estimate the epidemiologic and economic consequence of emergency vaccination, estimates of vaccine efficacy are important for simulation models.
- Several studies evaluated the efficacy of emergency vaccination. Variability between the estimates can be observed.

Objective

- Conduct a comprehensive assessment of the efficacy of emergency vaccination using meta-analysis of available literature.

What is meta-analysis?

- Statistical technique.
- Summarize the results of different studies that address a related hypothesis.
- Controls for study characteristics
- Results in an overall average are more powerful than outcomes of individual studies.

Characteristics of an emergency vaccine?

- Contain no residues of a live virus and have minimal side effects to newborns and adults
- After a single application with the recommended dose, achieve a potency dose 50 (PD50) of ≥ 6
- Be compatible with serological tests that identify infection in vaccinated animals
- Induce reasonably long lasting immunity and provide a broad spectrum of antigenic protection
- Be stable under storage once formulated
- Provide a rapid protection after vaccination
- Reduce the reproduction ratio (R_0) to below 1

Inclusion criteria

- Experimental challenge with FMD emergency vaccination using cattle, swine, and/or sheep.
- Research or symposium papers published in English language.
- Report the number of protected animals and the total number of animals in
 - vaccinated group
 - non-vaccinated control group.
- Both groups should be challenged with a homologous virus.
- Several groups were contacted to obtain unpublished data.

Definition of efficacy

- Determine the efficacy of FMD emergency vaccination in terms of:
 - Clinical protection.
 - Virological protection.

Outcome parameters

- Parameters to represent protection:
 - Relative risk (**RR**) of clinical disease.
 - RR of infection.

RR of clinical disease

- RR = the incidence of clinical disease in the vaccinated group divided by the incidence in the non-vaccinated group.
- Incidence of clinical disease = number of clinically diseased animals divided by the total number of animals per group.

Infection

- Several tests were used to diagnose infection.
 - When an animal was positive to any of the tests, the animal was considered infected.
- The tests were:
 - Virus isolation from the blood, oral, nasal, or esophageal-pharyngeal fluid.
 - Presence of antibodies to non-structural proteins.
 - RT-PCR of oral, nasal, or esophageal-pharyngeal fluid.

RR of virological protection

- The RR of virological protection = the incidence of FMD infection in the vaccinated group divided by the incidence in the non-vaccinated group.
- The incidence = the number of FMD infected animals divided by the total number of animals per group.

Meta-analysis procedure

- A pooled RR was calculated for the clinical and virological protection for
 - Cattle
 - Swine
 - Sheep
- A random effect model was used to calculate the pooled RR.
 - This model corrects for the variability between studies due to known and unknown covariates.
- An analysis was conducted when at least 4 studies were available.

Publication bias

- Studies with large and interesting effects → more likely to be published
- Studies with small or no effects → not likely to be published




- Consequence: biased body of research.
- Tests to detect publication bias:
 - a measures of study size relative to the study effect.


Results - descriptive

- 28 studies fitted the criteria.
- 27 studies included in the analyses.
- 10 conducted using cattle.
- 9 conducted using swine.
- 5 conducted using sheep.
- 3 had more than 1 species.
- 4 unpublished studies conducted on swine.


Results – Clinical protection in cattle



Results – Clinical protection in cattle - publication bias




Results – Clinical protection in cattle - publication bias




Results – Virological protection in cattle

- Vaccinated cattle
 - 0.65 (0.53 – 0.81) lower chance of FMD infection
- No significant publication bias was identified.

Results – Clinical protection in swine



Results – Virological protection in swine



Results – Protection in sheep

- Vaccinated sheep had
 - 0.31 (0.18-0.53) lower chance to develop clinical disease and
 - 0.57 (0.39-0.82) lower chance of FMD infection
- No significant publication bias was identified in both analyses.

Conclusions

- Emergency vaccination provided clinical and virological protection against FMD in cattle, swine and sheep.
- No significant publication bias was identified in the analyses.
- The outcomes of the meta-analysis can be used for further economic assessment of emergency vaccination in case of an outbreak using simulation modeling.

FLAVYDOL